

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number
WO 01/36681 A2

(51) International Patent Classification⁷: C12Q 1/68

STRANGE, Christina, M.; 6400 C Weatherby Ct., Frederick, MD 21703 (US). WILLIAMS, Inna, R.; 504 Carr Avenue, Rockville, MD 20850 (US). LORINCZ, Attila, T.; 6 China Berry Court, North Potomac, MD 20878 (US).

(21) International Application Number: PCT/US00/31277

(22) International Filing Date:
14 November 2000 (14.11.2000)

(74) Agents: MOROZ, Eugene et al.; Morgan & Finnegan, L.L.P., 345 Park Avenue, New York, NY 10154 (US).

(25) Filing Language: English

(81) Designated States (national): AU, BR, CA, JP.

(26) Publication Language: English

(30) Priority Data:
09/440,419 15 November 1999 (15.11.1999) US
09/707,178 6 November 2000 (06.11.2000) US

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(71) Applicant: DIGENE CORPORATION [US/US]; 1201 Clopper Road, Gaithersburg, MD 20878 (US).


Published:

— Without international search report and to be republished upon receipt of that report.

(72) Inventors: LAZAR, James, G.; 9914 Bellhaven Road, Bethesda, MD 20817 (US). ZAKEL, Joan, M.; 13317 Hunt Ridge Road, Ellicott City, MD 21042 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IMMUNOLOGICAL DETECTION OF RNA:DNA HYBRIDS ON MICROARRAYS



(57) Abstract: The present invention provides a hybridization method and kit for the detection and measurement of biological molecules. A test sample containing the biological molecules of interest is hybridized with an unlabeled or optionally a detectably labeled complementary biomolecule to form a double-stranded hybrid immobilized to a solid phase. The immobilized hybrid may be detected with an entity which specifically recognizes an RNA:DNA hybrid, followed by analyses and quantification. Therefore, the present invention provides a method and kit to detect and measure biological molecules that is simple to use, highly specific, sensitive, and accurate for screening a plurality of biological molecules.

WO 01/36681 A2

IMMUNOLOGICAL DETECTION OF RNA:DNA HYBRIDS

ON MICROARRAYS

FIELD OF THE INVENTION

5 The present invention is in the general field of detection of biological molecules, including DNA, RNA, protein and the like, and specifically in the field of detection of RNA:DNA hybrids on a solid phase, as further described herein, using a hybridization assay.

BACKGROUND OF THE INVENTION

10 The RNA or DNA for many genes, including those associated with disease states, and microorganisms and viruses have been isolated and sequenced. Nucleic acid probes based on such sequences are currently available to identify a large
15 number of genes and infections. Nucleic acid probes are detectable nucleic acid sequences that hybridize to complementary RNA or DNA sequences in a test sample. Detection of the probe indicates the presence of a particular nucleic acid sequence in the test sample for which the probe is specific. In addition to aiding scientific research,
20 nucleic acid probes may be used to detect the presence of viruses and microorganisms such as bacteria, yeast and protozoa as well as genetic mutations linked to specific disorders in patient samples.

 Grunstein, et al, Proc. Natl. Acad. Sci. USA 72:3961 (1975) and
25 Southern, J. Mol. Biol. 98:503 (1975) describe hybridization techniques using radiolabeled nucleic acid probes. Nucleic acid hybridization probes have the advantages of high sensitivity and specificity over other detection methods and do not require a viable organism. Hybridization probes are often labeled with a radioactive substance that may be easily detected.

30 The existing hybridization techniques that utilize radioisotopes to label probes introduce additional expenses caused by the high costs of disposal of radioactive waste products and the need for monitoring personnel and the workplace for contamination. In addition, the short half-life of radioactive compounds such as ³²P

0 requires that radioactive probes be produced frequently. Radioactive nucleic acid hybridization is therefore discouraged in commercial areas such as clinical diagnosis.

Probes have been indirectly labeled in an attempt to avoid the problems associated with direct radioactive labeling. One common method of indirect labeling
5 is to attach biotin, a small vitamin, to the nucleic acid probe using a chemical or enzyme technique. Following hybridization to the specific nucleic acid, the biotin is detected by reaction with streptavidin, a protein that binds biotin tightly and has been labeled with an enzyme or fluorochrome. Bound biotin-streptavidin complex may be
10 detected by reaction with color-producing substrates and the fluorochrome may be seen when reacted with incident light of appropriate wavelength. However, indirect labeling of hybridization probes with biotin or other haptens often increases the "hydrophobicity" of the probe. The probe tends to interact non-specifically with materials other than the complementary nucleic acid target, leading to high
15 background. The biotin label increases non-specific binding, which leads to high background, thereby reducing sensitivity and increasing the likelihood of a false-positive result. Indirect labeling is also less sensitive than direct labeling because the labeling density is limited; only a small fraction of the bases are labeled giving a
20 limiting number of sites for signal generation. An increase in the labeling density of a probe leads to increased non-specific binding, higher background, and ultimately, failure of the probe to hybridize with its target due to the interference of the hapten with base pairing. Indirectly labeled probes are therefore not well suited to clinical
25 diagnosis because of its inaccuracy and false positive results.

Hybridization of a probe to the specific nucleic acid sequences has been detected with the use of an intercalating agent such as acridine orange or ethidium bromide as described in U.S. Patent No. 4,563,417 to Albarella et al. The
30 intercalating agent becomes inserted between hybridized base pairs of probe and sample nucleic acids and causes the tertiary structure of the helix to unwind. An antibody specific for the newly formed antigenic determinant created by the intercalating agent and the unwound helix is detected by conventional means. This
35 method lacks selectivity for the target hybrids because intercalating agents fail to

9 ~~recognize specific sequences. Furthermore, the antibodies recognize only the~~
intercalating agent/ nucleic acid complex, but do not detect a specific sequence.
Therefore, additional selection or purification steps are required to prevent non-
specific signal, making this time consuming and labor intensive approach poorly suited
5 for clinical diagnosis

Hybridization of the probe to the specific nucleic acid sequences may
also be detected with the aid of an antibody specific for a labeled probe as described in
U.S. Patent No. 4,743,535 to Carrico. The probe is labeled with a detectable
10 substance such as flavin adenine dinucleotide (FAD) or a fluorescent agent. An
antibody specific for the labeled probe, after it has hybridized to the specific nucleic
acid sequence, is detected by a biochemical reaction. This method of detection also
creates non-specific binding and the likelihood of false-positive results and is not well
suited for clinical screening.

15 Attempts have been made to increase the sensitivity of nucleic acid
assays by target amplification. Methods of amplifying nucleic acid sequences are
commercially available. These methods include the polymerase chain reaction (PCR),
the ligation amplification reaction (LCR), and the transcription based amplification
20 reaction (TMA). PCR technology is described in PCR Protocols A Guide to Methods
and Applications by Michael A. Innis, David H. Gelfand, John J. Sninsky and Thomas
J. White, pp. 39-45 and 337-385 (Academic Press, Inc., Harcourt Brace Jovanovich,
Publishers, 1990). PCR technology is also described by Marx, J.L., Science 140:1408-
25 1410 (1988) and in U.S. Patent Nos. 4,683,195 and 4,683,202, to Mullis. Ligation
amplification reaction is described by Wu, D.Y and Wallace, R.B, Genomics 4:560-
569 (1989) and Barringer, K.J., et al., Gene 89:117-122 (1990). Transcription based
amplification reaction is described by Kwoh, D.Y., et al., Proc. Natl. Acad. Sci. USA
30 86:1173-1177 (1989). These methods have the advantages of high sensitivity, but the
disadvantages of having a lengthy, tedious, and expensive sample preparation, being
prone to false-positive results from reaction product contamination, and having the
inability to accurately quantify the initial amount of target nucleic acids.

35 Amplification reaction products are most often detected by a hybridization assay.

o

The degree of sensitivity achieved in assays for the detection of nucleic acid molecules, either RNA or DNA, in a sample is generally lower for RNA than DNA because RNA is subject to degradation by endogenous RNAses in the sample, resulting in less RNA available for detection. In addition, background interference
5 caused by contaminants in the sample is difficult to eliminate without causing further degradation of the target nucleic acid, such as RNA.

Hybridization assays for the detection of nucleic acid molecules, i.e. RNA, have been developed. For example, a hybridization protection assay for RNA is
10 commercially available from Gen-Probe Inc. (San Diego, CA). The hybridization protection assay employs a single-stranded nucleic acid probe linked to an acridinium ester, as described by Engleberg, N.C., ASM News 57:183-186 (1991), Arnold et al. Clin. Chem. 35:1588-1594 (1989) and U.S. Patent No. 4,851,330. Hybridization of
15 the probe to a target RNA molecule protects the acridinium ester bond from heat hydrolysis so that the detected chemiluminescent signal is proportional to the amount of target RNA in the sample. The sensitivity of this protection assay is limited by background luminescence caused by non-hybridized probe.

Polyclonal and monoclonal antibodies and other similar entities are
20 commonly used for detection purposes. Specifically, polyclonal antibodies recognize a plurality of epitopes, while monoclonal antibodies only recognize one specific epitope. Monoclonal antibodies which detect RNA:DNA hybrids are currently available. Polyclonal antibodies which detect RNA:DNA hybrids have been prepared, although,
25 generally, they have not been as specific as the monoclonal antibodies, which are designed to bind to a specific epitope.

Monoclonal antibodies to RNA:DNA hybrids are now available. U.S. Patent No. 4,732,847 to Stuart et al. and the publication of Stuart et al., Proc. Natl. Acad. Sci. USA 78:3751 (1981) describe a method of hybridization detection of
30 specific nucleic acid sequences on a solid surface involving a monoclonal antibody specific for a poly(A)-poly(dT) duplex. In Stuart, annealing DNA or RNA sequences complementary to the sequence of interest forms RNA:DNA hybrids. Stuart
35 specifically teaches against the use of polyclonal antibodies because with polyclonal

antibodies, one cannot preclude significant binding to single- or double- stranded nucleic acids. Further, unlike the present invention described herein, Stuart does not contemplate the advantages of polyclonal antibodies for arrays of very short oligomers on glass or silicon chips. In addition, Stuart does not contemplate microarrays, especially high-density arrays on glass slides or silicon chips. Nor does Stuart disclose attaching a nucleic acid probe to the surface of a solid phase. Instead, Stuart fixes a sample polynucleotide to a surface, while probe (e.g., a predetermined nucleotide sequence) is present in the liquid phase. In view of the foregoing, the present invention provides significant benefits and advantages to the art.

Boguslawski et al., J. Immunol. Methods 89:123-130 (1986) developed a hybridization assay using anti-hybrid coated polystyrene beads isolated on filter paper in an attempt to reduce non-specific binding and avoid complicated washing procedures. A monoclonal antibody specific for RNA:DNA hybrids secreted by hybridoma HB 8730, is disclosed in U.S. Patent No. 4,833,084 to Carrico et al. In Carrico, RNA:DNA hybrids formed by specific reannealing of a probe polynucleotide and the sequence of interest can be sensitively and specifically detected by binding to the monoclonal antibodies.

Microarrays refer to an orderly arrangement of distinct biological molecules, including RNA, DNA, protein, or the like, arrayed or immobilized to a solid substrate. These microarrays of binding agents, such as oligonucleotides and probes, have become an increasingly important tool in the biotechnology industry and related fields. Microarrays comprising a plurality of binding agents or elements are immobilized onto the surface of a solid support in an orderly fashion or pattern, find use in a variety of applications, including drug screening, nucleic acid sequencing, mutation analysis, and the like. Elements as used herein in a microarray context, refer to hybridizable nucleic acid sequences, oligonucleotides, primers, probes, and/ or amino acid sequences arranged in a distinct and identifiable manner on the surface of a substrate. Detection of biological molecules through the use of microarrays is beneficial for analyzing numerous samples and biological molecules, reducing the amount of sample required for analysis, decreasing experimental variability,

0
decreasing sample preparation time, confirming results, and for decreasing costs of such analysis.

Currently, one of the primary uses of microarrays is to measure gene expression in biological samples. Gene expression measurements include detecting the presence or absence of mRNA or measuring increased or decreased concentrations of mRNA. In order to detect hybridization and to measure gene expression by conventional methods, however, the sample must first be purified and labeled. Two common techniques for purifying and labeling the sample are: 1) RNA amplification, labeling, and hybridization, and 2) cDNA labeling and hybridization. The amplification part of the first technique is described in U.S. Patents 5,716,785 and 5,891,636 issued in 1998 and 1999, respectively, to Van Gelder et al. Highly purified total RNA or mRNA is used, which is an expensive and tedious time-consuming procedure. An oligo-dT primer is also used to reverse-transcribe the poly A-tailed mRNA into an anti-sense single-stranded cDNA. The oligo-dT further contains the sequence for T7 RNA polymerase on the 5 prime end of the dT sequences. After reverse transcription, a combination of RNase H, DNA ligase, and DNA polymerase are used to generate a double stranded cDNA. Because the original RT primer contained a T7 RNA polymerase promoter, the double-stranded cDNA contains a full T7 RNA promoter. The double-stranded cDNA is then used as a template for T7 RNA polymerase. Approximately 100-1000 additional copies of RNA are generated from each copy of cDNA. During the transcription process, labeled nucleotides are incorporated into the transcribed RNA. Labeled RNA is then hybridized to the DNA microarray forming labeled RNA: DNA hybrids. Fluorescent labels may be detected directly while indirect labels may be detected after reaction with a secondary binding agent.

30 A second sample preparation technique produces and measures labeled cDNA. In this technique total RNA or mRNA is purified from the biological sample. An oligo-dT primer is used to reverse-transcribe the poly-A tailed mRNA into an anti-sense single-stranded cDNA. During the reverse-transcription, labeled nucleotides are incorporated into the nascent DNA strand. After synthesis, the RNA strand is

35

destroyed. The labeled cDNA strand is then hybridized to the microarray. If the nucleotides were labeled with fluorescence, then the hybrids are visualized directly with a fluorescence array scanner. If the nucleotides were labeled with biotin, then the microarray is first reacted with labeled streptavidin and then scanned.

5 The disadvantages of both of these techniques are several fold. Firstly, both require a large quantity of highly purified nucleic acids (i.e. RNA or DNA). Purification requires additional steps which are time consuming and labor intensive. In addition, these techniques are inaccurate. Reverse transcription occurs at different
10 efficiencies and kinetic rates depending on the nucleic acid sequences, artificially changing the concentration of specific nucleic acid sequences. Prokaryotic mRNA and some eukaryotic mRNA do not contain the poly A sequence or tail at the 3 prime end or the poly A tail may be degraded during purification, and therefore cannot be labeled or detected with the current techniques since there is no sequence to prime the reverse
15 transcriptase step. The current techniques are thus restrictive to the types of samples which can be used for detection. Also, these methodologies involve labeled nucleotides. The incorporation of labeled nucleotides into unlabeled nucleic acids occurs at a lower efficiency and at a slower rate than natural nucleotides. Once more,
20 labels may be incorporated with different efficiencies depending on the sequence. Therefore, the label density may differ between different sequences; artificially changing the measured amount of these nucleic acids. Thus, quantification is only relative. Labeled nucleic acids also exhibit different hybridization kinetics than natural
25 nucleic acids, usually rendering them less specific. In addition, the present methods may require higher stringency hybridization conditions than unmodified nucleotides to achieve the same level of specificity. However, use of the higher stringency conditions to achieve acceptable specificity will lower the sensitivity of detection. Consequently,
30 there is a need for an assay for detection and for quantitative analysis of biological molecules, including DNA, RNA, protein, and the like, that is accurate, both time and cost efficient, and capable of screening one or more sample biological molecules with great sensitivity and minimal non-specific binding.

Therefore, it may be useful to have a method to detect and measure the amount of one or more biological molecules, including, but not limited to RNA, DNA, or protein, that is easy to use, highly specific, accurate, and sensitive for screening biological molecules.

5 Accordingly, it is an object of the invention to provide an assay to detect the absence or presence, and quantify biological molecules, including, but not limited to RNA, DNA, or protein.

It is also an object of the present invention to provide a method of
10 detecting an RNA:DNA hybrid comprising a specific target first biological molecule in a sample and a second biological probe.

It is an object of the present invention to provide a sensitive and quantitative assay having minimal false positives.

15 It is a further object of the present invention to provide an assay for massive parallel screening.

SUMMARY OF THE INVENTION

Disclosed is an assay for detecting and measuring a biological molecule
20 of interest, including RNA, DNA, protein, and the like, in a sample by hybridizing the biomolecule to a complementary biomolecule probe forming double-stranded hybrids, followed by immunological detection of these double-stranded hybrids formed on a solid phase with an antibody or other entity which specifically recognizes RNA:DNA
25 hybrids and is detectable. This method may be used to detect the presence of one or more specific biological molecules present in a variety of samples.

This invention provides for a method of simultaneously monitoring the amount (e.g. detecting and quantifying the amount) of a multiplicity of biological molecules.

30 The present invention relates to an assay for detecting RNA:DNA hybrids using detectably labeled entities specific for recognizing RNA:DNA hybrids. Preferably, the entity is a detectably labeled RNA:DNA hybrid-specific antibody or a fragment thereof. The antibody used for detecting the RNA:DNA hybrids may be
35

monoclonal or polyclonal, and preferably polyclonal for detection of short biological molecule probes having a length of less than 30 bases.

The present invention also relates to an assay using the microarrays of the invention to determine physiological responses by gene expression, polymorphism
5 mutation detection, SNP analysis, or the like. The method may be used to detect any and all genotypic variations, including insertion or deletion mutations.

Further, the present invention relates to an assay utilizing reverse transcriptase for extending short biological molecules, thereby enhancing detection of
10 RNA:DNA hybrids. Preferably, the reverse transcriptase is thermostable and lacks RNase H function.

The present invention further relates to a kit for the detection and quantification of biological molecules, wherein the kit may be used to screen samples
15 for large numbers of targets described herein by the present invention.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-D are a schematic representation of a preferred embodiment of the immunological antibody detection of RNA:DNA hybrids on
20 microarrays. Figure 1A shows hybridization of the RNA sample to the complementary DNA sequence that is attached to the microarray forming an RNA:DNA hybrid as depicted in Figure 1B. Subsequently, antibodies, either monoclonal or polyclonal, bind the RNA:DNA hybrids, as seen in Figure 1C. Figure 1D illustrates the detection
25 of fluorescent labels using a fluorescent laser scanner.

Figures 2A-D are a schematic representation of a second embodiment of the immunological detection of RNA:DNA hybrids on microarrays wherein the microarray comprises universal capture sequences. Figure 2A shows the hybridization
30 of the universal array with single-stranded DNA and sample RNA. Each hybrid formed on the microarray comprising a DNA:DNA region and an RNA:DNA region, as depicted in Figure 2B. Antibodies detect and bind RNA:DNA hybrids in Figure 2C. Figure 2D illustrates one means of detection comprising fluorescent antibody labels
35 using a fluorescent laser scanner.

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224

Figures 6A-B are graphs representing a solid phase-bound

oligonucleotide length comparison using detection with a monoclonal antibody (Figure 6A) and a polyclonal antibody (Figure 6B) as a function of signal to noise ratio of the microarray.

5

DETAILED DESCRIPTION OF THE INVENTION

An assay and kit are provided for the detection and quantification of one or more target biological molecule in one or more samples. In general, a test sample comprising biological molecules, including, but not limited to RNA, DNA,
10 protein, or the like, is collected and is either directly or indirectly, hybridized to a solid phase bound-nucleic acid probe specific for the target biomolecule. Non-hybridized nucleic acid sequences are removed, preferably by washing. Hybridization is then detected by a reaction with an RNA:DNA hybrid antibody that is labeled directly or
15 indirectly with a detectable label, and/ or detected by a labeled nucleic acid sequence which is complementary to the bound nucleic acid probe sequence.

In one embodiment of the present invention, a specific nucleic acid of a sample is hybridized to a complementary nucleic acid probe, preferably using an
20 oligonucleotide or other nucleic acid, which is spotted or synthesized to a solid phase, thereby forming a double-stranded RNA:DNA hybrid. Any entity which specifically recognizes RNA:DNA hybrids, preferably an antibody specific to an RNA:DNA hybrid, or fragment thereof, may be used for detection and measurement.

Also, the present invention utilizes short biological molecules,
25 preferably primers or probes, immobilized to a solid phase. It may be desirable to extend the primers with a reverse transcriptase, preferably one lacking RNase H function, enabling RNA:DNA hybrid-specific antibodies, RNA:DNA hybrid antibody fragments, or entities which specifically associate with RNA:DNA hybrids, to more
30 efficiently bind and be detected.

A further embodiment of the present invention encompasses three biological molecules, all of which are preferably nucleic acids. A first sample biomolecule hybridizes to a complementary second biological molecule, preferably a
35

0 probe, and either simultaneously or sequentially, hybridizes the second nucleic acid to a third nucleic acid, wherein one of the nucleic acids is immobilized to a solid phase, and the RNA:DNA hybrids which are formed, are detected by an entity specific for RNA:DNA hybrids.

5 In a further embodiment of the present invention, an immobilized biological molecule preferably comprising a protein, may bind to a sample biomolecule, preferably a nucleic acid, such that if the nucleic acid is an RNA:DNA hybrid, then it may be detected by an entity specific for RNA:DNA hybrids. For
10 example, DNA may bind a DNA binding site of the immobilized protein, wherein the DNA portion of the protein-DNA complex may be hybridized to RNA. In a similar manner, the immobilized protein may bind RNA, wherein the RNA portion of the protein-RNA complex may be hybridized to DNA. The resulting RNA:DNA hybrids
15 may be detected by an entity specific for RNA:DNA hybrids, such as an RNA:DNA hybrid-specific antibody or fragment thereof.

The present invention provides significant advantages to the art in its use of microarrays. Since either crude or purified sample may be used, the invention has a simplified sample preparation process, allowing for a more accurate detection
20 and measurement of biological molecules. Also, biological molecules need not be directly labeled for detection and measurement, thereby avoiding any interference attributed to the label. The present invention provides an extremely sensitive method for detecting and measuring biological molecules, since a very high labeling density
25 may be achieved by utilizing an entity that binds to RNA:DNA hybrids. Such exquisite sensitivity reduces the amount of sample required for analysis. Unlike other methods, the current invention may measure prokaryotic mRNA and some eukaryotic mRNA that lacks a poly A tail or has been degraded after purification.

30 Another advantage of the present invention is that reverse transcription is not required, but it may be employed if desired for enhanced sensitivity. One of the most advantageous aspects of the present invention is direct quantification of biological molecules. Unlike the commonly used techniques which only relatively
35 quantify RNA, e.g. 2-color competitive methods, the present invention utilizes a direct

~~approach to interpret results and a simplified analysis of biological molecules.~~ In addition, the present invention may simultaneously analyze a plurality of biological molecules due to its simplified sample process. Therefore the present invention allows much more straightforward interpretation and simplification of results.

5 In the present invention, a "probe" or a "nucleic acid probe", as used herein, is defined to be a collection of one or more nucleic acid or nucleic acid-like fragments whose hybridization to a second nucleic acid may be detected. The probe may be unlabeled or labeled as described below so that its binding to the second
10 nucleic acid may be detected. The probe may be produced from a source of nucleic acids from one or more particular portions of the genome, which may be known or unknown, for example one or more clones, an isolated whole chromosome or chromosome fragment, a collection of polymerase chain reaction (PCR) amplification
15 products, or a synthetic nucleic acid or PNA molecule. Alternatively, a probe may comprise a random, semi-random, or targeted sequence. The probe may be processed in some manner, for example, by blocking or removal of repetitive nucleic acids or enrichment with unique nucleic acids. Thus the word "probe" may be used herein to refer not only to the detectable nucleic acids, but to the detectable nucleic acids in the
20 form in which they are applied to the target, for example, with the blocking nucleic acids. The blocking nucleic acid may also be referred to separately. What "probe" refers to specifically is clear from the context in which the word is used. A probe may also function as a primer in the context of its use as an initiation point for
25 polymerization, i.e. for transcription or replication.

 The probe may also be isolated nucleic acids immobilized on a solid surface. In some embodiments, the probe may be a member of a microarray of nucleic acids as described, for instance, in WO 96/17958. Techniques capable of producing
30 high density microarrays may also be used for this purpose (see, e.g., Fodor et al. Science 767-773 (1991) and U.S. Pat. No. 5,143,854 to Pirrung, M.C.). Probes may also be deposited as elements onto the reaction substrate for interrogating the target molecules, and may be either directly or indirectly labeled.

0
The disclosed assay of the present invention may be used to detect and quantify any biological molecule, or combination of biological molecules in a sample, wherein the term "biological molecule" and "biomolecule" used interchangeably, as defined herein, refers to nucleic acids, amino acids, analogues, peptides, antibodies, and the like. "Nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof, from any source, including, but not limited to synthetic or derived from bacteria, yeast, viruses, and the cells or tissues of higher organisms such as plants or animals, and unless otherwise limited, may encompass known analogs of natural nucleotides that may function in a similar manner as naturally occurring nucleotides. Peptide nucleic acids (PNAs) are also encompassed within the scope of the term nucleic acid.

15 A "nucleic acid" is further defined herein as a single- or double-stranded nucleic acid ranging in length from 2 to about 10, 000 bases. As also used herein, the term "nucleic acid" refers to oligonucleotides, cDNA, mRNA, amplicons, plasmids, and the like. An "oligonucleotide" is one preferred nucleic acid probe comprising of at least 6 to about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20 to 25 nucleotides, which may be used in PCR amplification or a hybridization assay, or a microarray. As used herein, oligonucleotide is substantially equivalent to the terms "amplimers" and "oligomers", as commonly defined in the art, and may be used as "primers" and "probes" as described herein.

25 Also, unless otherwise limited, the term encompasses nucleic acids containing known analogues of natural nucleotides which have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. In addition, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g. degenerate codon substitutions) and complementary sequences as well as the sequence explicitly indicated.

35 Nucleic acid sequences for detection, referred to herein as nucleic acid molecules of interest, or target nucleic acid molecules, are selected based on the needs

0
and purpose of the detection. In general, a nucleic acid molecule of interest may be
chosen based on known criteria for selecting a nucleic acid sequence for detection. For
example, a particular nucleic acid molecule may be associated with a pathogen, a
disease state, or a predisposition to a disease, and detection of such a nucleic acid
5 molecule may have a diagnostic value. For example, mRNA specific to tumor cells or
normal cells may be detected. In addition, the disclosed method also allows the
detection of a biological molecules comprising, but not limited to, protein, peptides,
primers, and DNA or RNA molecules, generated by other biochemical or chemical
10 methods such as those generated by CAR, NASBA, etc. The detection of nucleic acids
also includes that of mutations, deletions, insertions of single nucleotide
polymorphisms, and other polymorphisms.

A "sample" or "target sample" as used interchangeably herein, is
15 defined in its broadest sense and includes both biological material and synthetic
material of biological molecules, including, but not limited to nucleic acids, amino
acids, proteins, peptides, and the like, and refers to a sample comprising total genomic
DNA, total RNA, genomic DNA or mRNA from, for example chromosomes, or
selected sequences (e.g. particular promoters, genes, amplification or restriction
20 fragments, cDNA, etc.) within particular amplicons or deletions. An embodiment of
the present invention is to detect either the presence or absence of the target nucleic
acid sample and to measure the amount of the sample that is to be quantified. The
term "target nucleic acid" may refer to the specific subsequence of a larger nucleic acid
25 to which the probe is directed to or to the overall sequence (e.g., gene or mRNA)
whose level is desired to detect, quantify, and determine the presence or absence. The
difference in usage will be apparent from the context.

The biomolecule sample may be extracted from particular cells or
30 tissues. The tissue sample from which the biomolecule sample is prepared is typically
taken from a patient suspected of having the disease associated with the amplification
or deletion being detected. In some cases, the biological molecules, for example,
nucleic acids, may be amplified using standard techniques such as PCR, prior to the
35 hybridization. The particular usage of the term "nucleic acid sample" will be readily

apparent to one of skill in the art from the context in which the term is used. For instance, the nucleic acid sample may be a tissue extract or cell lysate sample prepared by methods known in the art. The sample is prepared such that biological molecules of interest are released from cells and are available for hybridization.

5 Alternatively, a sample for the disclosed method of the invention may be from any source containing or suspected of containing nucleic acid. The source of nucleic acid may be in purified or non-purified form. Preferred types of samples, or sources of samples, that are suitable for use in the disclosed method are those samples
10 already known or identified as samples suitable for use in other methods of nucleic acid detection. Many such samples are known. For example, the sample may be from an agricultural or food product, or may be a human or veterinary clinical specimen. Samples may be a biological fluid such as plasma, serum, blood, urine, sputum, cell
15 lysate, or the like. The sample may contain bacteria, yeast, viruses and the cells or tissues of higher organisms such as plants or animals, suspected of harboring a biological molecule of interest. Methods for the extraction and/or purification of nucleic acids, for example, RNA have been described by Maniatis et al., *Molecular Cloning: A Laboratory Manual* (New York, Cold Spring Harbor Laboratory, 1982).

20 Since samples may also be in a crude or unpurified state, the sample preparation or processing is simplified. By using samples found in a more natural state, accurate expression detection and quantification is achieved. In addition, unlike other techniques which require the presence of a poly A sequence for priming the
25 reverse transcriptase step in order to label and detect sample, the present invention may be used to measure prokaryotic mRNA and eukaryotic mRNA that does not have a poly A tail at the 3 prime end.

 Target biological molecules of interest for use in the disclosed method
30 may come from various sources, both natural and synthetic. For example, various types of RNA include messenger RNA, ribosomal RNA, nucleolar RNA, transfer RNA, viral RNA and heterogeneous nuclear RNA, total genomic DNA, cDNA, proteins, peptides, or the like. In addition, whole naturally occurring entities or
35 fragments thereof may be used.

o

Solid phases or solid supports include, but are not limited to, those made of plastics, resins, polysaccharides, silica or silica-based materials, functionalized glass, modified silicon, carbon, metals, inorganic glasses, membranes, nylon, natural fibers such as silk, wool and cotton, and polymers. Solid phases or solid supports may be porous or non-porous. In some embodiments, the material comprising the solid support has reactive groups such as carboxy, amino, hydroxy, etc., which are used for covalent or non-covalent attachment of the probes. Suitable polymers may include, but are not limited to, polystyrene, polyethylene glycol tetraphthalate, polyvinyl acetate, polyvinyl chloride, polyvinyl pyrrolidone, polyacrylonitrile, polymethyl methacrylate, polytetrafluoroethylene, butyl rubber, styrenebutadiene rubber, natural rubber, polyethylene, polypropylene, (poly)tetrafluoroethylene, (poly)vinylidene fluoride, polycarbonate and polymethylpentene. Preferred polymers include those outlined in U.S. Pat. No. 5,427,779 to Elsner, H. et al., hereby expressly incorporated by reference. Solid phases and solid supports include, and are not limited to, any solid material to which the probes, primers, oligonucleotides, proteins, peptides, or the like, may be coupled or adhered. Solid phases and solid supports may have any useful form including thin films or membranes, beads, bottles, microwell plates, dishes, slides, fibers, woven fibers, shaped polymers, particles, chips and microparticles. Preferred substrate forms for a solid phase are microtiter dishes, silicon chips, glass slides, and tagged beads.

For general application, where a molecule is to be covalently bonded to the solid substrate surface, the surface may be activated using a variety of functionalities for reaction, depending on the nature of the bound component and the nature of the surface of the solid substrate. Thus the surface of the solid substrate, if required, may be modified by the introduction of functionalities which may then react with the bound component.

"Microarrays" comprise a plurality of different biological molecules including cDNA, amplicons, plasmids, proteins, peptides, and the like, wherein plurality encompasses at least two different biological molecules, wherein the biomolecules are immobilized to a solid phase in an ordered matrix or structure. In

theory, there need be only one component, but in a preferred embodiment there will be at least 10, more usually at least 20, frequently at least 50, desirably 100 or more, and even 1,000 or more, but usually not more than about 10^4 , more usually not more than about 100,000, with from about 10 to 10,000 immobilized to a solid phase or solid support being preferred. While theoretically the number of different components may exceed 10^5 , due to the ability to specifically have a small amount or volume at a specified finite site, for the most part there is no need to exceed 100,000 and such large numbers of different components do add some complexity to the preparation of the microarray. As the number of components immobilized to a solid phase will usually not exceed 10^5 , the number of individual addressable sites may be substantially larger, depending on the nature of the bound component, the source of the signal, the nature of the signal which is detected, the sensitivity with which the signal may be detected, the nature of the bound microarray, such as the size of the microarray, the manner in which the microarray is produced, and the like. Therefore, microarrays are preferably used for "massive parallel screening", described herein as the simultaneous screening of at least about 10, preferably about 1,000, and more preferably about 10,000, different biological molecule hybridizations.

One preferred form of a microarray comprises a spotted array to which 1-10, 10-100, or most preferably more than 100 separate nucleic acids, preferably oligonucleotides, primers, or the like, may be deposited, may be spotted or synthesized as an array of small dots or elements, as described herein. These nucleic acids, deposited, spotted, or synthesized on a solid phase, are referred to herein as "elements". Typically, an element will be less than about 1 mm in diameter. Generally, element sizes are from 1 μ m to about 5 mm, preferably between about 1 μ m and about 1 mm. Nucleic acid primers for use in the disclosed method may be synthesized using established oligonucleotide synthesis methods. Such methods range from standard enzymatic digestion followed by nucleotide fragment isolation (see for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edition (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) Chapters 5, 6) to purely synthetic methods, for example, by the cyanoethyl phosphoramidite method

using a Milligen or Beckman System I-Plus DNA synthesizer (for example, Model 8700 automated synthesizer of Milligen-Bioscience, Burlington, MA or ABI Model 380B). Synthetic methods useful for making oligonucleotides are also described by Ikuta et al. (Ann. Rev. Biochem. 53:323-356 (1984), (phosphotriester and phosphite-triester methods)), and Narang et al. (Methods Enzymol., 65:610-620 (1980), (phosphotriester method)).

Another form of microarray is a three dimensional array, examples of which include an array of color-coded beads (Luminex; Austin, TX) and an array of radiofrequency-tagged beads (PharmaSeq; Monmouth Junction, NJ). A three dimensional microarray, as used herein, is any solid phase having three dimensions, wherein each microarray comprises a plurality of different biological molecules, preferably nucleic acid primers, attached to the surface. Thus, the location of each primer on the solid phase microarray enables the identification of each nucleic acid primer sequence. Manipulations of the disclosed assay may be utilized. For example, a three-dimensional microarray comprising of a plurality of nucleic acid primers may be mixed with target nucleic acids of interest. When the primers are short, it may be desirable to extend these molecules with polymerases, such as for example reverse transcriptase, so as to incorporate the binding capacity of the RNA:DNA hybrid-specific entity, as described herein, including antibodies and fragments thereof. By capturing the antibodies on a solid phase, the primers of the solid phase microarray on which RNA:DNA hybrids have formed may be separated from the primers where no hybrid has formed. The entities specific for RNA:DNA hybrids may then be detected and the identities of the primers determined. Many other assay schemes may be used for the disclosed method.

The microarray has emerged as a preferred format for the miniaturization of assays that detect and measure RNA, DNA, proteins, and the like, for application towards, for example, gene expression, mutation and polymorphism analysis, SNPs, detection of genetic variations, etc. Microarrays allow the level of tens to several thousands of genes or genetic variations (for example SNPs) to be measured from a single sample on a single device. A weakness of the traditional microarray

0
methods is that the biological molecule, preferably nucleic acid (either RNA or DNA) to be measured, must first be labeled, often through conversion of one type of nucleic acid to another, for example RNA to labeled DNA, so that it may be detected and measured.

5 The present invention preferably utilizes a "nucleic acid microarray", which as defined herein, comprises a plurality of nucleic acid sequences, including, but not limited to, DNA, RNA, amplicons, plasmids, and the like, immobilized to a solid support to which complementary target nucleic acids are hybridized. The nucleic acids
10 of the microarray may, for example, contain sequence from specific genes or clones, probes, primers, or oligonucleotides, bound to a porous or non-porous solid phase or solid support. Nucleic acids of various dimensions may be used in the microarrays of the invention.

15 The nucleic acids may be coupled to the solid support or substrate. Such a microarray is a solid support to which multiple different nucleic acids have been coupled or adhered in an array, grid, or other organized pattern. "Nucleic acid microarrays" preferably comprise arrays of nucleic acid sequence strands on silicon chips, glass slides, or other solid support, and are in widespread use for detection and
20 measurement of gene expression, mutation and polymorphism analysis, etc. Several methods are available for preparing nucleic acid microarrays. Strands of nucleic acid sequences may be non-covalently or covalently bound to a solid substrate through passive or chemical coupling methods. Other approaches utilize synthetic methods to
25 build the nucleic acid molecules directly on the surface of the substrate. A simpler, but more limited approach, is to prepare labeled nucleic acid sequences and then bind the labeled nucleic acid sequences to a substrate that has been coated with a binding partner.

30 Alternatively, elements of proteins and/ or peptides may be coupled to the solid support or substrate in an organized pattern. These immobilized protein elements may be bound by nucleic acids, proteins, peptides, and/ or nucleic acid hybrids. Detection is achieved using entities specific for RNA:DNA hybrids, such as
35 antibodies or fragments thereof. If the protein or peptide binds RNA:DNA hybrids,

0 then the RNA:DNA hybrid portion of the protein-hybrid complex may be detected using an entity specific for RNA:DNA hybrids. If the protein binds DNA, then the DNA portion of the protein-DNA complex can be hybridized to RNA resulting in the formation of an RNA:DNA hybrid. If the protein binds RNA, then the RNA portion of
5 the protein-RNA complex may be hybridized to DNA, resulting in the formation of an RNA:DNA hybrid. The RNA:DNA hybrids may be detected using an entity specific for RNA:DNA hybrids, such as RNA:DNA hybrid-specific antibodies, or their fragments thereof.

10 A "hybrid" is a double-stranded nucleic acid comprising RNA or DNA. The duplex may be DNA:DNA, RNA:RNA, or RNA:DNA, or may comprise artificial nucleotides. An RNA homoduplex is a base-paired double-stranded RNA. An RNA:DNA heteroduplex comprises an RNA strand and a strand comprising DNA
15 nucleotide monomers. All or a region of the duplex may be double-stranded. Typically, at least 10 bases of the duplex will be double-stranded. The phrases "to specifically hybridize" or "specific hybridization" or "selectively hybridize to", or the like, refer to the binding, duplexing, or hybridizing of a nucleic acid molecule preferentially to a particular nucleotide sequence under stringent conditions when that
20 sequence is present in a complex mixture (e.g., total cellular) DNA or RNA.

Nucleic acid probes immobilized on a solid substrate allow formation of RNA:DNA hybrids localized on the substrate. Such localization provides a convenient means of washing away reaction components that might interfere with subsequent
25 detection steps, and a convenient way of assaying for multiple different target nucleic acid sequences simultaneously. RNA:DNA hybrids may be independently formed at each site where a different primer is adhered. For immobilization of probes to form a solid phase microarray of biological molecules, the methods described herein may be
30 used.

An "entity", as defined herein, refers to any molecule which specifically recognizes RNA:DNA hybrids. Examples of entities that may recognize RNA:DNA hybrids may include, but are not limited to, chimeric antibodies, and natural or

0 genetically-engineered proteins or nucleic acids that specifically bind to RNA:DNA hybrids.

One preferred embodiment of entity is "antibody". As used herein, antibody is intended to be used in the broadest sense and to include whole, intact
5 antibodies, antibody fragments, recombinant antibodies, chimeric antibodies, polyfunctional antibody aggregates, or in general any antibody-derived substance that comprises at least one antibody combining site having the characteristics described herein or other entities. Preferably, in the present invention, these entities, specifically
10 detect and bind RNA:DNA hybrids. Antibodies of any of the known classes and subclasses of immunoglobulins are contemplated, for example, IgG, IgM, and so forth, as well as active fragments such as the IgG fragments conventionally known as Fab, F(ab'), and F(ab')₂. Antibodies may comprise monoclonal antibodies (including
15 agonist, antagonist, and neutralizing antibodies) which bind to a specific epitope and polyclonal antibodies having polyepitopic specificity, or other entities.

Any antibodies or entities specific for double-stranded RNA:DNA hybrids may be used to directly detect the hybrid of the invention. In the present invention, polyclonal antibodies are preferred in the embodiment which utilizes them
20 for detecting short nucleic acid sequences, preferably those less than 30 bases in length.

The antibodies used to detect RNA:DNA hybrids may be either monoclonal or polyclonal antibodies. It may also be advantageous to use a mixture of
25 monoclonal and polyclonal antibodies. Furthermore, the invention includes the use of customized polyclonal or monoclonal antibodies that may be produced with specific binding properties. For instance, monoclonal or polyclonal antibodies that specifically bind to very short (less than 20 base pairs) RNA:DNA hybrids may be produced and
30 may find use in detecting very short RNA:DNA hybrids. In addition, monoclonal or polyclonal antibodies may be produced that are either more or less sensitive to mismatches within the RNA:DNA hybrid. Antibodies which are more sensitive to mismatches within the RNA:DNA hybrid will find extra utility in the detection of
35 genetic variation while antibodies which are less sensitive to mismatches with the

RNA:DNA hybrid will find use in the detection and quantification of specific classes of nucleic acids. Other antibodies may also be used that specifically detect nucleic acid triplexes (DNA: RNA:DNA or RNA:DNA:RNA) or DNA:PNA or RNA:PNA hybrids, wherein PNA is defined herein as peptide nucleic acid.

5 Polyclonal antibodies directed against the RNA:DNA hybrids are prepared by injecting a suitable laboratory animal with an effective amount of the peptides or antigenic component, collecting serum from the animal, and isolating specific sera by any of the known immunoadsorbent techniques. Animals which may
10 readily be used for producing polyclonal RNA:DNA hybrid antibodies include chickens, mice, rabbits, rats, goats, horses, and the like. In a preferred embodiment of the present assay, a polyclonal RNA:DNA hybrid antibody is derived from goats immunized with an RNA:DNA hybrid. Hybrid-specific antibody is purified from the goat serum by affinity purification against RNA:DNA hybrid immobilized on a solid
15 support.

Monoclonal antibodies, prepared by standard techniques, may be used in place of the polyclonal antibodies. A variety of techniques may be used to obtain suitable antibodies specific for RNA:DNA hybrids. (For example, U.S. Patent
20 Number 4,833,084 to Carrico, U.S. Patent Number 4,732,847 to Stuart et al. and Stuart et al., Proc. Natl. Acad. Sci. USA 78:3751 (1981)). A monoclonal antibody specific for RNA:DNA hybrids, secreted by hybridoma HB 8730, is disclosed in U.S. Patent No. 4,833,084 to Carrico. Preferably, in accordance with the present invention,
25 monoclonal antibodies are used for the detection of nucleic acids greater than 30 bases in length.

The isolation of anti-RNA:DNA hybridomas has improved the development of assays for genetic mutations linked to specific defects and the
30 detection of bacterial and viral infections. However, assays utilizing these RNA:DNA hybrid-specific monoclonal antibodies often suffer from a high level of non-specific binding causing false positive results. Boguslawski et al., J. Immunol. Methods 89:123-130 (1986) developed a hybridization assay using anti-hybrid coated
35

polystyrene beads isolated on filter paper in an attempt to reduce non-specific binding and avoid complicated washing procedures.

The preferred antibody for RNA:DNA hybrids is prepared by the method of Kitawaga, Y. and Stollar, B.D., Mol. Immunology 19:413-420 (1982) or
5 according to the method set forth in U.S. Patent No. 4,732,847, issued March 22, 1988 to Stuart et al., both of which are incorporated herein by reference.

The identification of the presence of the hybrids may be achieved by employing either polyclonal or monoclonal antibodies or other entities specific for the
10 RNA:DNA hybrid complex. Detection may be achieved by labeling either the antibody specific for the hybrid RNA:DNA complex, or by employing labeled antibodies which bind to the anticomplex. For example, where the antibody is derived from a mouse, antibodies to mouse antibodies, for example rabbit anti (-mouse IgG),
15 may be labeled so as to bind to any anticomplex bound to the complex bound to the solid support.

A wide variety of labels have been used in other environments which may be applicable here. One of the more common labels is radionuclides, which may be used with autoradiography to visualize the areas of binding. Another label is a
20 fluorescer such as fluorescein, mercocyanine, or rhodamine which by irradiation with light of excitation, the presence of fluorescence may be monitored. Alternatively, an enzyme may be used which results in a product which may be detected and localized in the area of the enzyme. A large number of dyes or metals capable of reduction may be
25 employed to provide detection. Common enzymes include horseradish peroxidase, glucose oxidase, galactosidase, alkaline phosphatase, or the like. The particular label or manner in which the detectable signal is observed is not critical to this invention. By employing antibodies to the anticomplex, the number of labels associated with a
30 particular binding of the anticomplex to the complex may be greatly amplified.

To facilitate detection of resulting binding of the antibody, or the other entity specific for double-stranded hybrids, to the hybrid, the antibody will normally be labeled with a detectable chemical group. Examples of detectable chemical groups
35 that may serve as labels are enzymatically active groups, such as coenzymes, enzyme

substrates, enzyme inhibitors, and enzymes themselves, fluorescers, chromophores, luminescers, specifically bindable ligands such as biotin or haptens which are detectable by binding of labeled avidin or labeled hapten antibodies, and radioisotopes.

In order for complete hybridization to occur, the optimal conditions are necessary for forming double-stranded hybrids. The term "stringent conditions" refers to conditions under which a probe will hybridize preferentially to a complementary sequence, and to a lesser extent to, or not at all to, other sequences. Complementarity between two single-stranded molecules may be "partial", in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands and in the design and use of PNA molecules. A "stringent hybridization" and "stringent hybridization wash conditions" in the context of nucleic acid hybridization experiments, such as, for example, Southern and Northern hybridizations are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes part 1 chapter 2. "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, N.Y.

"Bind(s) substantially" refers to complementary hybridization between a probe nucleic acid and a target nucleic acid and embraces minor mismatches that may be accommodated by reducing the stringency of the hybridization media to achieve the desired detection of the target polynucleotide sequence hybridized to the bound oligonucleotide sequence, which includes cDNA, amplicons, plasmids, and the like.

Hybridization of the probe nucleic acid to the nucleic acid molecule of interest may be carried out under any suitable conditions, and preferably under conditions which favor hybridization and form double-stranded hybrids. See for

0
example, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edition
(Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989).

For example, in one embodiment of the present invention, a primer is
needed to begin reverse transcription. A "primer" is defined herein, as a nucleic acid
5 molecule that may anneal to a DNA or RNA template molecule and serves as the
initiation point for nucleic acid synthesis. A custom primer is generally a synthetic
oligonucleotide, including cDNA, amplicons, plasmids, and the like, but naturally
occurring nucleotides act as primers as well, both *in vitro* and *in vivo*. *In vitro* uses of
10 primers include, for example, cDNA synthesis, Sanger dideoxy sequencing, and PCR.

This particular embodiment requires a nucleic acid "template" of
interest in order to identify the target nucleic acid(s) of interest once the target nucleic
acid sample(s) are obtained. Wherein, "nucleic acid template" or "template" as used
interchangeably herein, is defined as a polynucleotide sequence from which
15 information is read to direct synthesis of another macromolecule. For example, this
may refer to a DNA strand being copied during DNA synthesis or transcription of
RNA, to an RNA strand being copied during reverse translation.

A primer of the disclosed method may be an oligonucleotide, cDNA,
20 amplicons, plasmids, and the like, either RNA or DNA, having sequence
complementary to a region on a nucleic acid molecule of interest. As used herein, the
complementary sequence of the primer is referred to as the "complementary portion".
As used herein, the region on the target nucleic acid molecule of interest
25 complementary to the primer is referred to as the "primer complement region". The
primer complement region of a target nucleic acid molecule of interest may be any
region of the target molecule of interest. For the embodiment of the present assay
which utilizes reverse transcriptase, a preferred mode comprises the primer
30 complement region of a target nucleic acid molecule be at some distance from the 5
prime end of the template nucleic acid molecule. This provides a longer region of
nucleic acid template between the site of primer hybridization and the end of the
template nucleic acid molecule, thereby amplifying the amount of RNA:DNA hybrid
35 to be detected.

o
In general, the primer complement region of a nucleic acid molecule of interest is chosen based on known criteria for selecting a nucleic acid sequence for detection. For example, to detect a particular nucleic acid molecule from among other nucleic acid molecules, it is preferred that the primer complement region is
5 characteristic of, or unique to, the target nucleic acid molecule of interest. If it is desired that any of a class of RNA molecules be detected, it is preferred that the primer complement region is chosen to have a sequence that is the same or substantially the same in all of the target nucleic acid molecules of interest. Once a primer complement
10 region is selected, the sequence of the primer is designed or chosen to be complementary to the chosen primer complement region of the molecule of interest. Any nucleic acid molecule for which a sequence is known or for which a sequence may be derived may be detected using the disclosed method.

15 In the method of the invention the complementary portion of a primer has a length that supports specific and stable hybridization between the primer and the primer complement region. Generally a primer of the present invention comprises 10 to 100 nucleotides, but is preferably 15 to 30 nucleotides.

20 The ability to characterize an individual by its genome is due to the inherent variability of genetic information. Although DNA sequences which code for necessary proteins are well conserved across a species, there are regions of DNA which are non-coding or code for portions of proteins which do not have critical functions and therefore, absolute conservation of nucleic acid sequence is not strongly
25 selected for. These variable regions are identified by genetic markers. Typically, genetic markers are bound by probes such as oligonucleotides or amplicons which specifically bind to unique variable regions of the genome. In some instances, the presence or absence of binding to a genetic marker identifies individuals by their
30 unique nucleic acid sequence. In other instances, a marker binds to nucleic acid sequences of all individuals but the individual is identified by the position in the genome bound by a marker probe. The major causes of genetic variability are addition, deletion or point mutations, recombination and transposable elements within
35 the genome of individuals in a plant population. The present invention may be applied

to detecting and measuring genotypic variation. For example, polymorphisms, such as SNPs, which are represented by different sequences, may be detected.

In general, the present invention assay involves the following steps:

1. Preparing a biomolecule probe or microarray of probes bound to a solid substrate (such as, for example, on plates, slides, wells, dishes, beads, particles, cups, strands, chips, and strips, both porous and non-porous) by spotting or synthesizing the biomolecule probe to a solid phase through standard chemical techniques;
2. Adding the target sample containing the first biological molecule of interest to the immobilized second biomolecule probes and allowing RNA:DNA hybrids to form;
3. Adding a detectable entity specific for RNA:DNA hybrids (including RNA:DNA hybrid-specific antibodies or fragments thereof); and
4. Detecting the entity bound to the immobilized RNA:DNA hybrids.

Another embodiment of the present invention involves the following steps:

1. Preparing a biomolecule probe or microarray of probes bound to a solid substrate (such as, for example, on plates, slides, wells, dishes, beads, particles, cups, strands, chips, and strips, both porous and non-porous) by spotting or synthesizing the biomolecule probe to a solid phase through standard chemical techniques;
2. Adding the target sample containing the first biological molecule of interest to the immobilized second biomolecule probes and allowing RNA:DNA hybrids to form;
3. Adding reverse transcriptase, preferably lacking RNase H function and thermostable.
4. Incubating under conditions that promote reverse transcription which extends the sequence, thus forming a much longer RNA:DNA hybrid and enhancing antibody detection.
5. Adding a detectable entity specific for RNA:DNA hybrids (including RNA:DNA hybrid-specific antibodies or fragments thereof); and
6. Detecting the entity bound to the immobilized RNA:DNA hybrids.

~~A further embodiment of the present invention includes the following~~

steps:

1. Preparing a biomolecule probe or microarray of probes bound to a solid substrate (such as, for example, on plates, slides, wells, dishes, beads, particles, cups, strands, chips, and strips, both porous and non-porous) by spotting or synthesizing the biomolecule probe to a solid phase through standard chemical techniques;
2. Adding the target sample containing the first biological molecule of interest to a second biomolecule probe bound to a microarray and a third unbound biomolecule probe;
3. Hybridizing the first target biological molecule to a complementary region of the third biomolecule probe;
4. Hybridizing the immobilized second biomolecule probe to an unhybridized complementary region of the third biomolecule probe;
5. Adding a detectable entity specific for RNA:DNA hybrids (including RNA:DNA hybrid-specific antibodies or fragments thereof); and
6. Detecting the entity bound to the immobilized RNA:DNA hybrids.

Another embodiment of the present invention includes the following steps:

1. Preparing a biomolecule probe or microarray of probes bound to a solid substrate (such as, for example, on plates, slides, wells, dishes, beads, particles, cups, strands, chips, and strips, both porous and non-porous) by spotting or synthesizing the biomolecule probe to a solid phase through standard chemical techniques;
2. Adding the target sample containing the first biological molecule of interest to a second biomolecule probe bound to a microarray and a third unbound detectably-labeled biomolecule probe;
3. Hybridizing the first target biological molecule to a complementary region of the second solid phase-bound biomolecule probe and forming an RNA:DNA hybrid;
4. Hybridizing the solid phase-bound second biomolecule probe to a complementary region of the third detectably labeled biomolecule probe;

5. Adding a detectable entity specific for RNA:DNA hybrids (including RNA:DNA hybrid-specific antibodies or fragments thereof); and

6. Separately detecting both the entity specific for RNA:DNA hybrids bound to the immobilized RNA:DNA hybrids and the detectably labeled biomolecule probe.

5 The disclosed assay may be used to detect a plurality of different biological molecules of interest in a sample. This is preferably accomplished by either screening for a sequence that is present in each of the target biological molecules of interest, or by screening with multiple probes that are collectively complementary to
10 regions on the biological molecules of interest. The latter approach is preferred for use in detecting, for example, some diseases or predispositions to disease that are associated with numerous different mutations to particular genes, or genetic variations, including, but not limited to insertion or deletion mutations. The present invention
15 also provides an assay which may be applied to a variety of applications, including, but not limited to gene expression, biological molecule (i.e. RNA, DNA, protein) detection on microarrays, mutation and polymorphism detection (i.e. SNP), and the like. In one particular embodiment, it is preferred to screen for sequences that are complementary
20 to the regions of the mutant nucleic acid products of these genes that are characteristic of each of the mutations. Thus, one major advantage of this assay is the high-throughput application, enabling large screenings of a plurality of samples and potential diseases.

 The disclosed method may also be used to determine the ratio of
25 expression of different biological molecule species from individual organisms or an individual sample. For this purpose, the method is used to detect multiple species simultaneously. Microarray detection, as disclosed herein, is useful for this purpose. The disclosed method may also be used to detect similar or related biomolecule
30 sequences where the related biological molecules have a common sequence motif between them, but which are otherwise different. For example, cells may contain multiple biological molecule species having similar regulatory sequences, similar structural motifs, or other sequences in common. Such classes of nucleic acid

⁰
~~molecules may be detected with a single probe species by designing the probe to~~
hybridize to the common sequence.

In the disclosed assay, an entity specific for RNA:DNA hybrids, including RNA:DNA hybrid-specific antibodies and their fragments, is utilized to
5 detect biological molecules that have hybridized to the probe microarray rendering the labeling of the target biomolecules no longer necessary, but an option. In this approach, the longer the RNA:DNA hybrid, the greater the signal since a longer RNA:DNA hybrid may bind more antibody than a short RNA:DNA hybrid.

10 Therefore, the longer the nucleic acid probe strands on the microarray, the more sensitive the detection of target nucleic acids or alternatively, the greater the signal intensity for a given amount of hybridized target nucleic acids. Unfortunately, it becomes more difficult and increasingly expensive to synthesize, prepare or utilize
15 longer strands of probes in the preparation of these microarrays.

One disclosed embodiment of the present assay describes relatively short nucleic acid probe sequences bound to a solid substrate, minimizing the time, effort, and expense necessary to create the microarray. Target nucleic acid sequences in the sample are hybridized to these short probes creating a short RNA:DNA hybrid
20 with a long nucleic acid tail. This short RNA:DNA hybrid probably only binds 1 or 2 RNA:DNA antibodies. When reverse transcriptase is added, and conditions are such that reverse transcriptase occurs, the nucleic acid probe portion of the RNA:DNA hybrid is extended to the length of the target nucleic acid strand, thus greatly
25 increasing the length of the RNA:DNA hybrid. If the target nucleic acid strand were 1500 bases in length, then the resulting RNA:DNA hybrid would approach 1500 base pairs. An RNA:DNA hybrid of this length binds significantly more RNA:DNA antibodies, thereby greatly increasing the intensity of the signal produced, and
30 increasing the sensitivity of detection of specific target nucleic acid sequences

One disclosed embodiment is a method of detecting target nucleic acid sequences by reverse transcribing all or part of the bound nucleic acid probe sequence with a reverse transcriptase lacking an RNA:DNA hybrid-dependent exonuclease
35 function (commonly referred to as an RNase H function or component) and detecting

- 32 -

the resulting RNA:DNA hybrid with an antibody specific for RNA:DNA hybrids. The nucleic acid probes are immobilized on a solid support in order to associate the RNA:DNA hybrid with the solid support. This allows for easy separation of hybrids from sample solution and specific detection of nucleic acid molecules based on the position of the hybrid on the solid support.

In one method of the present invention, reverse transcription is carried out using a reverse transcriptase, preferably a reverse transcriptase lacking RNase H function. The reaction mixture including the nucleic acid molecule of interest, preferably in this embodiment, RNA; the hybridized immobilized nucleic acid primer; and the reverse transcriptase is then incubated under conditions to allow reverse transcription of the RNA molecule of interest and formation of DNA:RNA hybrids. Examples of reverse transcriptases that may be used in the disclosed method, or that may be adapted for use in the disclosed method are listed in Table 1. Preferred reverse transcriptases for use in the present method include reverse transcriptases, 18053-017, 18064-014 and 18064-071 from Life Technology; reverse transcriptases M5301 and M5302 from Promega; and reverse transcriptase 600085 from Strategene; each disclosed in Table 1.

Table 1

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
	1 unit incorporates 1 nmol TTP into acid-insoluble form/10 min at 35°C using poly(A).oligo dT ₁₂₋₂₈ as substrate	0.2 M KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable RNase, exonuclease	Adv Biotech AB-0321 AB-0321b
26,700 U/mL	1 unit incorporates 1 nmol dTMP into acid-precipitable form/10 min at pH 8.3, 37°C	200 mM KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable endonuclease, RNase	ACS Heideib F00750S F00750M
10-20 U/μl	1 unit incorporates 1.0 nmol [³ H]-JTTP into acid-insoluble products/10 min at 37°C	0.2 M KPO ₄ , 2.0 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable endonuclease, exonuclease, RNase	Amersham E 70041Y E 70041Z
>50,000 U/mg;	1 unit incorporates 1 nmol [³ H]-dTMP into acid-precipitable products/10 min at 37°C using poly(A).d[pT] ₁₅ as template primer	200 mM KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable nonspecific RNases, nonspecific DNases (gel electrophoresis)	Boehringer 10911B 149506Z
30,000 u/mL	1 unit incorporates 1 nmol [³ H]-TMP into nucleic acid product/10 min at 37°C	Solution containing 0.2 M KPO ₄ , 2.0 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable nonspecific nuclease	ICN 855928 855929

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
13 U/ μ L	1 unit incorporates 1 nmol dTNP into a TCA-insoluble product/10 min at pH 8.3, 37°C	0.2 M KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable RNase, exogenous, nicking or degradation of RNA	NBL Gene 020704
30 U/ μ L	1 unit incorporates 1 nmol dTNP into a TCA-insoluble product/10 min at pH 8.3, 37°C	0.2 M KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2; for high efficiency synthesis of full length cDNA in the 6-10 kilobase range	No detectable RNase, exonuclease, endonuclease, nicking	NBL Gene 020703
>20,000 U/mg; 10,000-20,000 U/mL	1 unit incorporates 1 nmol dNTP into DE-81 adsorbable form/10 min at 37°C	200 mM KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable RNase, DNase	Oncor 120111 120112
25,000-50,000 U/mg protein; 10,000-20,000 U/mL	1 unit incorporates 1 nmol dTMP into acid-insoluble product/10 min at pH 8.3, 37°C using poly(A)p[dT] ₁₂₋₁₈ as template primer	Molecular biology grade; homogeneous purity; solution containing 0.2 M KPO ₄ , 2.0 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable RNase, DNase, nickase	Pharmacia 27-0922-01 27-0922-02
10,000-70,000 U/mL	1 unit incorporates 1.0 nmol [³ H]-dTTP into acid-insoluble product/10 min at 37°C	Purified; 20 mM KPO ₄ , 2 mM DTT, 0.2% Triton X-100, 50% glycerol, pH 7.2	No detectable endonuclease, nonspecific RNase	Stratagene 600081 600082
>40,000 U/mg; >20,000 U/mL	1 unit incorporates 1 nmol TMP in acid-insoluble product/10 min at 37°C with poly(A)[dT] ₁₈ as substrate	Coomassie Blue shows a single band purity; 50 mM Tris-HCl, 10 mM DTT, 100 mM NaCl, 0.05% polydocanol, 1 mM EDTA, 50% glycerol, pH 8.4	No detectable nonspecific RNases, nonspecific DNases (gel electrophoresis)	Boehringer 1062603

5 10 15 20 25 30 35

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
50-250 U/ μ L	1 unit incorporates 1 nmol deoxynucleotides into acid-precipitable material/10 min at 37°C using poly(rA).oligo[dT] ₁₂₋₁₈ as template primer	>90% purity by SDS-PAGE; 20 mM Tris.HCl, 0.1 M NaCl, 0.1 mM EDTA, 1 mM DTT, 0.01% NP40, 50% glycerol, pH 7.5	No detectable non-specific ss- and ds-endonuclease, exonuclease, RNase	Amersham E 70456Y E 70456Z
50,000 U/mL	1 unit incorporates 1 nmol TTP into acid-insoluble form/10 min at pH 8.0, 37°C using poly(rA).oligo[dT] as template primer	50 mM Tris.HCl, 0.1 mM DTT, 100 mM NaCl, 1 mM EDTA, 0.1% NP40, 50% glycerol, pH 8.3	No detectable endonuclease, RNase	ACS Heideib F00755S F00755M
200 U/ μ L	1 unit incorporates 1 nmol deoxyribonucleotide into acid-precipitable material/10 min at pH 8.3, 37°C using poly(A)-oligo-[dT] ₁₂₋₁₈ as template primer	Purity by SDS-PAGE, 250 mM Tris.HCl, 15 mM MgCl ₂ , 375 mM KCl, pH 8.3 and 100 mM DTT	No detectable RNase H	Life Technol 18053-017
200 U/ μ L	1 unit incorporates 1 nmol deoxyribonucleotide into acid-precipitable material/10 min at pH 8.3, 37°C using poly(A)-oligo-[dT] ₁₂₋₁₈ as template primer	Purity by SDS-PAGE, 250 mM Tris.HCl, 15 mM MgCl ₂ , 375 mM KCl, pH 8.3 and 100 mM DTT	No detectable RNase H	Life Technol 18064-014 18064-071

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
200 U/ μ L	1 unit incorporates 1 nmol deoxyribonucleotide into acid-precipitable material/10 min at pH 8.3, 37°C using poly(A)-oligo [dT] ₁₂₋₁₈ as template primer			Life Technol 28025-013 28025-021
25,000 U/mL	1 unit incorporates 10 nmol TTP into acid-insoluble material/10 min at 37°C using poly(rA).oligo(dT) as template primer	0.1 mM NaCl, 50 mM Tris.HCl, 5 mM DTT, 1 mM EDTA, 0.1% NP40, 50% glycerol, pH 7.6	No detectable endonuclease, RNase	NE Biolabs 253S 253L
≥ 5000 U/mg protein	1 unit incorporates 1.0 nmol [³ H]-TMP into acid-insoluble products/10 min at 37°C using poly(A).d[pT] ¹⁸ as substrate	Recombinant; 99% by HPLC, SDS-PAGE; lyophilized containing 0.2% BSA as stabilizer	No detectable nuclease	Boehringer 1465333
20-40 U/ μ L	1 unit incorporates 1 nmol deoxyribonucleotide into DE-81 absorbable form/10 min at 37°C	Overproducer; 50 mM Tris.HCl, 0.1 M NaCl, 0.1% Triton X-100, 1 mM EDTA, 5 mM DTT, 50% glycerol, pH 8.3	No detectable endo and exodeoxyribonucleases, RNases	Fermentas EP0351 EP0352
100-200 U/ μ L	1 unit incorporates 1 nmol dTTP into acid-insoluble form/10 min at pH 8.3, 37°C	Recombinant; $\geq 90\%$ purity by SDS gel; 5X reaction buffer, 250 mM Tris.HCl, 375 mM KCl, 15 mM MgCl ₂ , 50 mM DTT, pH 8.3	No detectable RNase H <1% DNase <3% RNase $\geq 90\%$ supercoiled plasmid	Promega M5301 M5302

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
	1 unit incorporates 1 nmol TTP into acid-insoluble form/10 min at 37°C using poly(A)-oligo(dT) ₁₁₋₁₈ as substrate	50 mM Tris.HCl, 0.1 M NaCl, 1 mM EDTA, 5 mM DTT, 0.1% Triton X-100, 50% glycerol, pH 8.3	No detectable RNase, exonuclease	Adv Biotech AB-0322 AB-0322b
	1 unit incorporates 1 nmol dTTP into acid-insoluble form/10 min at 37°C	50 mM Tris.HCl, 5.0 mM DTT, 1.0 mM EDTA, 0.1 M NaCl, 0.1% NP40, 50% glycerol, pH 8.0	No detectable endonuclease, RNase	CHIMERx 1375-01 1375-02
35,000 U/mg	1 unit incorporates 10 nmol dTTP into acid-insoluble material/10 min at pH 8.6, 37°C using oligo(dT) ₁₂₋₁₈ primed poly(A) _n as template	Solution containing 50% glycerol, 50 mM Tris.HCl, 0.1 M NaCl, 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100, pH 7.5	No detectable RNase, endonuclease, exonucleolytic DNase, protease	Epicentre M4425H M4410H
	1 unit incorporates 1 nmol labeled dATP into acid-insoluble material/10 min at 37°C	Solution containing 0.1 mM NaCl, 50 mM Tris.HCl, 1 mM EDTA, 5 mM DTT, 0.1% NP40, 50% glycerol, pH 8.0		ICN 152020
	1 unit incorporates 1 nmol TMF into DE-81 adsorbable form/10 min at 37°C using poly(A)-oligo(dT) ₁₂₋₁₈ as substrate	50 mM Tris.HCl, 0.1 M NaCl, 5 mM DTT, 1 mM EDTA, 0.1% Triton X-100, 50% glycerol, pH 8.3	No detectable RNase, DNase	Oncor 120301 120302

5

10

15

20

25

30

35

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
50,000-95,000 U/mg protein;	1 unit incorporates 1 nmol dTMP into acid-insoluble product/10 min at pH 8.3, 37°C using	Molecular biology grade; homogeneous purity; solution containing 50 mM Tris.HCl, 0.1 M NaCl, 1 mM EDTA, 5 mM DTT, 0.1% Triton X-100, 50% glycerol, pH 8.3	No detectable RNase, DNase, nickase	Pharmacia 27-0925-01 27-0925-02
10,000-20,000 U/mL	poly(rA).p[dT] ₁₂₋₁₈ as template primer			
50,000 U/mL	1 unit incorporates 1.0 nmol [³ H]TTP into acid-insoluble product/10 min at 37°C	50 mM Tris.HCl, 5 mM DTT, 1 mM EDTA, 100 mM NaCl, 0.1% NP40, 50% glycerol, pH 8.0	No detectable RNase H, DNase, nonspecific RNase	Stratagene 600085
10-30 U/μL	1 unit incorporates 1 nmol [³ H]dTMP/10 min at 37°C with poly(rA).oligo(dT) as template primer	200 mM KPO ₄ , 2 mM DTT, 0.2% NP40, 50% glycerol, pH 7.2	No detectable non-specific nuclease	Amersham E 2610Y E 2610Z
400 U and 1600 U	1 unit incorporates 1 nmol [³ H]dTMP/10 min at pH 8.3, 37°C with poly(rA).oligo(dT) as template primer	Solution containing 200 mM KPO ₄ , 2 mM DTT, 0.2% NP40, 50% glycerol, pH 7.2	No detectable nuclease	Takara 2610

SPECIFIC ACTIVITY	UNITS DEFINITION	PREPARATION FORM	ADDITIONAL ACTIVITIES	SUPPLIER CATALOG NO.
	Transcriptase: 1 unit incorporates 4 nmol dTTP into acid insoluble material/30 min at pH 8.3, 45°C using oligo(dT)18-primed poly(A) _n as template; DNA Polymerase: 1 unit incorporates 10 nmol dNTP into acid insoluble material/30 min at pH 8.3, 74°C	Solution containing 50% glycerol, 50 mM Tris.HCl, 0.1 M NaCl, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween 20, 0.5% NP40, pH 7.5; no detectable DNA exo- and endonuclease, protease, RNase		Epicentre Retrotherm™ RT R19250 R19500 R1910H
	1 unit incorporates 1 nmol dTTP into acid-insoluble form/10 min at 50°C	50 mM Tris.HCl, 5.0 mM DTT, 0.1 mM EDTA, 50% glycerol, stabilizers, pH 7.5	No detectable endonuclease, 3'-exonuclease, 5'-exonuclease/5'-phosphatase, nonspecific RNase, ss- and ds-DNase	CHIMERX 1374-01 1374-02

0

5

10

15

20

25

30

35

Reverse transcription may generally be performed at any temperature within the functional temperature range of the reverse transcriptase. Preferably, the temperature of incubation is any temperature at which the reverse transcriptase is functional and the primer remains hybridized to the target nucleic acid molecule. For non-thermophilic reverse transcriptases, preferred temperatures are those temperatures that are at or around the optimum temperature for the reverse transcriptase. For most non-thermophilic reverse transcriptases this temperature will be between about 25°C and 45°C.

In a preferred embodiment, a thermophilic reverse transcriptase is used for increasing selectivity. The highest temperature at which a thermophilic reverse transcriptase is functional may be quite high. For this reason, preferred temperature ranges for reverse transcription when a thermophilic reverse transcriptase is used are most conveniently described in terms of the calculated melting temperature of a hybrid between the RNA molecule of interest and the primer. Such a melting temperature is referred to herein as the RNA/primer melting temperature (R/P T_m). Preferred ranges include a temperature from 20°C below the melting temperature of a hybrid between the RNA molecule of interest and the primer and 5°C above the melting temperature of a hybrid between the RNA molecule of interest and the primer. Other preferred ranges when using a thermophilic reverse transcriptase include those listed in Table 2.

TABLE 2:

MAX:	to R/P T _m	to 5°C below R/P T _m	to 3°C below R/P T _m
1	20°C below R/P T _m	20°C below R/P T _m	20°C below R/P T _m
2	15°C below R/P T _m	15°C below R/P T _m	15°C below R/P T _m
3	10°C below R/P T _m	10°C below R/P T _m	10°C below R/P T _m
4	7°C below R/P T _m	7°C below R/P T _m	7°C below R/P T _m
5	5°C below R/P T _m	5°C below R/P T _m	5°C below R/P T _m
6	3°C below R/P T _m	3°C below R/P T _m	3°C below R/P T _m

It is specifically noted that every specific, but unnamed, range within the enumerated ranges above is contemplated as an alternative preferred range. Preferred temperatures for reverse transcription include about 20°C below R/P T_m, about 15°C below R/P T_m, about 12°C below R/P T_m, about 10°C below R/P T_m, about 7°C below R/P T_m, about 5°C below R/P T_m, about 3°C below R/P T_m, 20°C

- 41 -

below R/P T_m, 15°C below R/P T_m, 12°C below R/P T_m, 10°C below R/P T_m, 7°C below R/P T_m, 5°C below R/P T_m, and 3°C below R/P T_m. In general, the closer the temperature is to the R/P T_m, the greater the degree of discrimination there will be between specific and non-specific hybrids of the RNA and primer. If the temperature is close to the R/P T_m, however, decreased stability of specific hybrids may cause priming to be less efficient.

R/P T_m may be determined either by calculation or by empirical measurement. For calculating R/P T_m, any established formula for calculating stability of nucleic acid hybrids may be used. A preferred formula for calculating R/P T_m is T_m =
$$\frac{\Delta H}{\Delta S + R \times \ln(C/4)} + 16.6 \log \frac{[K^+]}{1 + 0.7 [K^+]} - 237.15,$$

which was derived from studies on the stability of perfectly-matched DNA:DNA hybrids. For RNA:DNA hybrids, incorporating formamide concentration in the formula does not hold because the relationship between formamide concentration and the depression of T_m is not linear. At 80% formamide, RNA:DNA hybrids are more stable than DNA:DNA hybrids, increasing the T_m by about 10 to 30°C depending on the sequence (Hames & Higgins, Nucleic Acid Hybridisation: A Practical Approach (IRL Press Limited, Oxford, England. 1985)). Carrying out the reaction in 80% formamide may therefore also be used to suppress formation of DNA:DNA duplexes, to preferentially select RNA:DNA hybrids, and to estimate the T_m for R/P. Because the empirically derived formulas for the estimation of RNA:DNA hybrid T_m may not be as accurate for short nucleic acid primers, the hybridization temperature is preferably determined by assessing hybrid stability in 0.1 - 0.4 M monovalent cation at temperatures ranging from 40 to 60°C. R/P T_m may also be determined empirically (Lesnick and Freier, Biochemistry 34:10807-10815 (1995), McGraw et al., Biotechniques 8:674-678 (1990), and Rychlik et al., Nucleic Acids Res. 18:6409-6412 (1990)).

As used herein, a thermophilic reverse transcriptase is any reverse transcriptase that retains at least 5% of its maximum activity at any temperature above

50°C or which has an optimal temperature of at least 50°C. Preferred reverse transcriptases are those which have an optimal temperature of at least 50°C. As used herein, maximum activity of a reverse transcriptase is defined as the activity, as measured in the assay described below, that a given reverse transcriptase exhibits at its optimal temperature. As used herein, optimal temperature of a reverse transcriptase is defined as the temperature at which the activity of the reverse transcriptase is greatest, as measured in the assay described below. The optimal temperature for a given reverse transcriptase may be determined by measuring its activity in the following assay at various temperatures. In general, an optimal temperature need be determined only to within a range so that assays need only be performed at intervals of 5 to 10 degrees.

Methods for immobilization of nucleic acid sequences to solid phase substrates are well established. Oligonucleotides, including half probes and rolling circle replication primers, may be coupled to substrates using established coupling methods. For example, attachment methods are described by Pease et al., Proc. Natl. Acad. Sci. USA 91(11):5022-5026 (1994), and Khrapko et al., Mol. Biol (Mosk) (USSR) 25:718-730 (1991). A method for immobilization of 3'-amine oligonucleotides on casein-coated slides is described by Stimpson et al., Proc. Natl. Acad. Sci. USA 92:6379-6383 (1995). A preferred method of attaching oligonucleotides to solid-state substrates is described by Guo et al., Nucleic Acids Res. 22:5456-5465 (1994).

The immobilization and arraying of nucleic acids or primer molecules to solid supports may be accomplished using any suitable technique. For example, immobilization may be accomplished either by *in situ* nucleic acid synthesis (Maskos and Southern, Nucleic Acids Research, 20:1679-1684 (1992); Pease et al., Proc. Natl. Acad. Sci. USA, 91:5022-5026 (1994)) or by covalent or passive attachment of chemically synthesized oligonucleotides (Guo et al., Nucleic Acids Research, 22:5456-5465 (1994)), or by covalent or passive attachment of other nucleic acids, amplicons, cDNAs, and the like, in combination with robotics arraying technologies. Other immobilization techniques are described in U.S. Patent No. 5,412,087 to McGall et al., U.S. Patent No. 5,429,807 to Matson et al., and U.S. Patent No. 5,510,087 to Fodor et

al. Thousands of different primers may be arrayed onto a small area on a solid support to interrogate thousands of target nucleic acid molecules. The density of nucleic acids or primers should be matched with the method of arraying and the means of detection.

One embodiment of the present invention comprises hybridization of
5 target nucleic acid sequences to the universal array comprising specific nucleic acid probes, wherein a "universal array" or "universal array sequences", herein interchangeably defined as short nucleic acid sequences of every possible base combination. The universal array sequences comprise a range of 6 - 10 bases,
10 preferably 5 - 6 bases, wherein the number of possible combinations (i.e. different probes bound to the solid phase) is 1024 and 4096, respectively, and enable nucleic acid expression analysis, wherein the result may be used as a fingerprint, in which different tissues or samples give different fingerprints.

The embodiments comprising a third nucleic acid probe may be
15 immobilized to the solid phase array by using "capture tags". As used herein, a capture tag is any compound that may bind to another compound or moiety. The primer is thus immobilized through binding of an attached capture tag to its binding partner. Such binding partners are referred to herein as "capture docks". A capture tag
20 is a compound, such as a ligand or hapten, that binds to or interacts with another compound, such as ligand-binding molecules or an antibody. It is also preferred that such interaction between the capture tag and the capture dock be a specific interaction, such as between a hapten and an antibody or a ligand and a ligand-binding molecule.

25 A further embodiment of this assay comprises a capture tag with two adjacent regions: a target nucleic acid-specific region and a "capture sequence complement". As used herein, a "capture sequence complement" comprises nucleic acid sequence which is complementary to "universal capture sequences" immobilized
30 to the solid phase microarray, wherein "universal capture sequences" refer to short nucleic acid sequences which are known and their location on the solid phase microarray are predetermined. The capture tag or probe comprising a "capture sequence complement" may be immobilized to the solid phase microarray by
35 hybridizing to its complementary "universal capture sequence".

o

~~In another embodiment of the present invention, the "capture tag" refers~~
to a labeled nucleic acid probe which hybridizes to its "capture dock" which is bound
to the solid phase microarray, wherein the capture dock is a common sequence specific
for the labeled nucleic acid probe.

5 Alternative capture tags include hapten or ligand molecules that may be
coupled to oligonucleotides. Capture tags, described in the context of nucleic acid
probes, have been described by Syvnen et al., Nucleic Acids Res., 14:5037 (1986).
Capture tags also include biotin, which may be incorporated into nucleic acids.

10 Adhering or coupling primers to a substrate may be accomplished by
adhering or coupling capture docks to the substrate. The capture docks mediate
adherence of a primer by binding to, or interacting with, a capture tag on the primer.
Capture docks immobilized on a substrate allow capture of the primer on the substrate.
15 By attaching different capture docks to different regions of a substrate different capture
tags attached to different primers, may be captured at different, and therefore
diagnostic, locations on the substrate. For example, in a microtiter plate multiplex
assay, capture docks specific for up to 96 different capture tags may be immobilized on
a microtiter plate, each in a different well. Capture and detection will occur only in
20 those wells corresponding to capture tags for which the corresponding nucleic acid
molecules were present in a sample.

 In one embodiment, the capture dock is an oligonucleotide. Methods
for immobilizing and coupling oligonucleotides to substrates are well established. For
25 example, attachment methods are described by Pease et al., Proc. Natl. Acad. Sci.
USA 91(11):5022-5026 (1994), and Khrapko et al., Mol Biol (Mosk) (USSR) 25:718-
730 (1991). A method for immobilization of 3'-amine oligonucleotides on casein-
coated slides is described by Stimpson et al., Proc. Natl. Acad. Sci. USA 92:6379-
30 6383 (1995). Another method of attaching oligonucleotides to solid phase substrates is
described by Guo et al., Nucleic Acids Res. 22:5456-5465 (1994).

 Methods for immobilizing proteins to substrates are well established.
Immobilization may be accomplished by attachment, for example, to aminated
35 surfaces, carboxylated surfaces or hydroxylated surfaces using standard

immobilization chemistries. Examples of attachment agents are cyanogen bromide, succinimide, aldehydes, tosyl chloride, avidin-biotin, photocrosslinkable agents, epoxides, maleimides and glutaraldehyde. These and other attachment agents, as well as methods for their use in attachment, are described in Protein immobilization:

- 5 Fundamentals and Applications, Richard F. Taylor, ed. (M. Dekker, New York, 1991), Johnstone and Torpe, Immunochemistry In Practice (Blackwell Scientific Publications, Oxford, England, 1987) pages 209-216 and 241-242, and Immobilized Affinity Ligands, Craig T. Hermanson et al., eds. (Academic Press, New York, 1992).

- 10 Proteins may be attached to a substrate by chemically cross-linking a free amino group on the antibody to reactive side groups present within the substrate. For example, proteins may be chemically cross-linked to a substrate that contains free amino or carboxyl groups using glutaraldehyde or carbodiimides as cross-linker agents. In this method, aqueous solutions containing free proteins are incubated with the solid-state
15 substrate in the presence of glutaraldehyde or carbodiimide. Standard immobilization chemistries are known by those of skill in the art.

- In another embodiment, the sensitivity of the disclosed method is increased by repeated washing of the hybrid sample to remove free unhybridized
20 nucleic acids present in the sample. It is useful to remove the non-specific unhybridized nucleic acid because secondary structures in the nucleic acid may be recognized by the detection means, resulting in elevated assay background.

- The preferred hybridization sample nucleic acid detection kits for use
25 with the disclosed method may be made using some or all of the components required for the method. The kit preferably contains an immobilized primer that is complementary to a region on a nucleic acid molecule of interest, and more preferably contains a plurality of immobilized primers that are each complementary to a region on
30 a nucleic acid molecule of interest.

- Preferably kits contain all or some of the following components: a sample transport medium for stabilization of the sample; a solid phase bound
microarray of biomolecules specific for a second biomolecule to be detected;
35 hybridization buffer; entity specific for RNA:DNA hybrids; wash buffer; enhance

~~buffer, and the reagents necessary for detecting the RNA:DNA hybrid-specific~~

antibody. In addition, some kits may include a thermostable reverse transcriptase lacking RNA:DNA hybrid-dependent exonuclease (RNase H) function. A further composition of the kits may include a nucleic acid probe comprising a capture
5 sequence complement region. In addition, the kits may also include a labeled biomolecule probe which hybridizes to a common sequence of the solid phase bound biomolecule. Kits may further comprise, but is not limited to, a universal array of biological molecules for the detection of a sample biomolecule. Kits may comprise all
10 of these components, or a portion thereof.

For amplified-antibody detection, in addition to the reagents included in the hybridization kits for direct detection described above, the following reagents comprising all or part may also be included in the kits: detectably labeled anti-mouse
15 IgG; biotinylated anti-mouse IgG; labeled anti-mouse streptavidin; biotinylated anti-streptavidin; or acetylated BSA solution.

The kits should contain a negative control and a positive control. Preferably, probes for the negative and positive controls are included on the solid
20 phase with the nucleic acid sequences.

The following non-limiting examples illustrate use of the present assay and kits.

Examples

It is understood that the disclosed invention is not limited to the
25 particular methodology, protocols, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

30 It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a host cell" includes a plurality of such host cells, reference to "the antibody" is a reference to one or more antibodies and
35 equivalents thereof known to those skilled in the art, and so forth.

0
Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of skill in the art to which the disclosed invention belongs. Although any methods and materials similar or equivalent to those described herein may be used in the practice or testing of the present invention, the preferred methods, devices, and materials are as described. Publications cited herein and the material for which they are cited are specifically incorporated by reference. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

10 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

15 Example 1

Detection of RNA:DNA Hybrids on Probe Microarrays

The following is an example of a preferred method of performing one embodiment of the disclosed method for the detection of target biomolecule sequences in a sample.

20 In general, the assay is preferably used to detect a sample size of 0.05 μg of nucleic acids. Most preferably, 0.05 μg - 10 μg of total nucleic acids is detected using the assay of the present invention.

25 The target nucleic acid sample was resuspended in nuclease-free water and added to hybridization buffer. The hybridization solution was denatured at 95°C for 2-5 minutes. The hybridization solution containing the target nucleic acid was added to the glass slide which has a microarray of oligonucleotides spotted. Hybridize at 65°C for 16-20 hours. Follow by either direct detection or amplified detection.

30 For direct detection, the glass slide with a microarray of primers bound to the surface was washed 3 times for 1-2 minutes with 1XPBS/ 0.05% Tween 20™ and shaken on a rotary shaker (1100 rpm). The RNA:DNA antibody staining solution was added so that the final concentration was 0.144 $\mu\text{g}/\mu\text{l}$. The glass slide microarray was incubated in solution for 1 hour at room temperature shaking (1100 rpm). The

35

glass slide was washed with 1XPBS/ 0.05% Tween 20™ and shaken (1100 rpm) for 15 minutes at room temperature. The microarray was then incubated in a mouse antibody, specifically directed against Cy3 or Cy5, staining solution at room temperature and shaken for 1 hour at 1100 rpm. A number of fluorescent dyes may be used, such as, but not limited to Cy3 or Cy5. The final concentration of the Cy-dye was 0.04 µg/µl in a solution of 10% goat serum and 1XPBS/ 0.05% Tween 20™. Using slightly rigorous means, the slide was washed four times for approximately 10 seconds each in wash buffer. The slide was then incubated at 53°C for 15 minutes in enhance buffer. The slide was then washed in wash buffer four times for approximately 10 seconds each, using mildly rigorous means. The microarray bound to the glass slide was dried by centrifugation at 2000 rpm for 7-10 minutes, or until dry. The results were analyzed by reading the slide in an array scanner (Affymetrix 417 Array Scanner or equivalent) with photo excitation at 532 nm and 635 nm, for scanning slides developed with Cy3 and Cy5 labeled antibodies, respectively.

For amplified detection, the slide was washed 3 times for 1-2 minutes in 1XPBS/ 0.05% Tween 20™, shaken on a rotary shaker at 1100 rpm. The slide was incubated in RNA:DNA hybrid-specific antibody staining solution for 1 hour shaking (1100 rpm) at room temperature. The final concentration of RNA:DNA antibody staining solution was 0.144 µg/µl. The slide was washed in 1XPBS/ 0.05% Tween 20™ for 15 minutes at room temperature shaking (1100 rpm). The spotted slide was covered with biotinylated mouse IgG antibody from goat staining solution and incubated for 10 minutes at room temperature. The microarray was washed 2 times for 1-2 minutes each in 1XPBS/ 0.05% Tween 20™. The spotted slide was covered with goat anti-mouse R-phycoerythrin streptavidin (0.01 µg/µl; SA-PE) staining solution and incubated for 10 minutes at room temperature. Washing was repeated as described above. The spotted slide was covered with biotinylated goat antibody raised against streptavidin (0.5 mg/ mL) staining solution and incubated for 10 minutes at room temperature. The slide was again incubated with goat anti-mouse R-phycoerythrin streptavidin (0.01 µg/µl; SA-PE) staining solution for ten minutes at room temperature. The slide was then washed 3 times for 1-2 minutes in 1XPBS/ 0.05%

0 Tween-20™. The microarray was then dried by centrifugation at 2000 rpm for 7-10 minutes, or until dry. The results were analyzed by reading slide in an array scanner or equivalent (Affymetrix 417 Array Scanner), with photo excitation at 532 nm and 635 nm, for scanning slides developed with Cy3 or PE and Cy5 labeled antibodies
5 respectively.

Example 2

Labeled Oligonucleotide Hybridization Prior to Sample Hybridization

10 A Cy3 or Cy5 labeled n-mer oligonucleotide was added to a hybridization buffer containing SSC and SDS and was denatured at 95°C for 2-5 minutes. A number of fluorescent dyes may be used, such as, but not limited to Cy3 and Cy5. The glass slide comprising spotted oligonucleotides was covered with the
15 hybridization solution and incubated at room temperature for 20 seconds. The coverslip was removed with rigorous dunks in 2XSSC/ 0.2% SDS. Any residual SDS was washed off by dunking the slide in 0.05XSSC for 30 seconds. The slide was dried by centrifugation at 2000 rpm for 7-10 minutes or until dry. The glass slide was then
20 read in an array scanner (Affymetrix 417 Array Scanner or equivalent), with photo excitation at 532 nm and 635 nm, for scanning slides developed with Cy3 and Cy5 labeled oligonucleotides, respectively, for analysis. Sample hybridization followed.

Example 3

Detection of RNA Mediated by Reverse Transcriptase

Lacking RNase H Function

25 The following is an example of a method of performing one embodiment of the disclosed method for the detection of target nucleic acid sequences in a sample.
30

The 5 prime biotinylated 20 to 30 nucleotide primers was mixed with a streptavidin coated solid phase and incubated for 30 to 60 minutes at 20-27°C with constant shaking (1100 rpm). A sample of target nucleic acids was added to the solid phase. Hybridization/ extension buffer (100 mM Tris-HCl, pH 8.3, 150 mM KCl, 6
35

- 50 -

0 mM MgCl₂, 20 mM DTT, and 1 mM each dNTP) was then added. The target nucleic acid and primer were annealed by heating the mixture to the optimal annealing temperature, preferably 60°C (optimal annealing temperature varies with primer and nucleic acid utilized), for 20-30 minutes. The mixture was then cooled at 20-27°C for 5 10 minutes. Additional hybridization/ extension buffer and reverse transcriptase, preferably thermostable and lacking RNase H, was added. The reaction was incubated for 30-60 minutes at 42°C. EDTA (0.5 M) was added and incubated for 30 minutes at 37°C. RNA:DNA hybrid-specific alkaline phosphatase conjugated antibody mix was 10 added and incubated at 20-27°C for 30-60 minutes. Any unbound antibody was washed away, followed by the addition of a chemiluminescent substrate. The solution was incubated for 15-30 minutes at 20-27°C. The signal which was emitted was read with a luminometer at the appropriate wavelength.

15 Example 4

Binding of RNA:DNA Hybrid-specific Antibodies

Hybridized RNA:DNA samples were incubated with the antibodies for a sufficient amount of time to allow conjugation of the hybrids. The hybrids were 20 bound to the antibodies by incubation for 5 minutes to 24 hours at 15 to 65°C on a platform shaker with a shaking speed of 0 to 1500 rpm. Preferably, the incubation time was 30 to 120 minutes at 20 to 40°C, with shaking at 300 to 1200 rpm. Most preferably, binding occurred with incubation at one hour at room temperature with 25 vigorous shaking on a rotary platform shaker with a rotary shaking speed between approximately 300 and 1000 rpm. It will be understood by those skilled in the art that the incubation time, temperature, and shaking may be varied to achieve alternative capture kinetics as desired.

30 Example 5

Oligonucleotide Length Comparison Detected by Monoclonal and Polyclonal Antibody

A single oligonucleotide of varying length was spotted at four different 35 concentrations in replicates of ten. The spotted 72-mer oligonucleotide was part of

IMAGE clone #259983 which corresponds to the 40S Ribosomal protein S11. The shorter oligonucleotides were sequential truncations of the parent 72-mer. These microarrays were hybridized to varying concentrations of complementary RNA and were visualized using the monoclonal primary RNA:DNA hybrid antibody. At a target
5 concentration of 800 pM, substantial signal was observed at an oligonucleotide length of 30 bases, but there was a drop in signal at 25 bases. Figure 6A shows the results of the signal to noise ratio as a function of oligonucleotide length at an RNA concentration of 800 pM with various spotted oligonucleotide concentrations upon
10 RNA:DNA hybrid-specific monoclonal antibody detection.

The polyclonal antibody detection protocol was the same as that described above for the monoclonal antibody, with the exception that a Cy3-labeled goat secondary antibody from rabbit was used instead of the mouse antibody from goat. The results (Figure 6B) with the polyclonal antibody showed significantly
15 improved detection compared to the monoclonal antibody for oligonucleotides less than 30 bases in length. For oligonucleotides greater than 30 bases, there was no significant difference in signal.

The polyclonal RNA:DNA antibody provided a significantly more
20 sensitive method for detecting RNA:DNA hybrids that are less than 30 base pairs long as compared to detection with the monoclonal antibody (see Figs. 6A-B).

The contents of all patents, patent applications, published PCT applications and articles, books, references, reference manuals and abstracts cited
25 herein are hereby incorporated by reference in their entirety to more fully describe the state of the art to which the invention pertains.

As various changes may be made in the above-described subject matter without departing from the scope and spirit of the present invention, it is intended that
30 all subject matter contained in the above description, or defined in the appended claims, be interpreted as descriptive and illustrative of the present invention. Many modifications and variations of the present invention are possible in light of the above teachings.

CLAIMS**What is claimed:**

1. A microarray method for detecting an RNA:DNA hybrid comprising the steps of:
 - 5 a) hybridizing a first biological molecule of the sample to a second immobilized biological molecule to form a double-stranded hybrid; and
 - b) detecting the RNA:DNA hybrid using a detectable entity which specifically binds to a component of the RNA:DNA hybrid.
- 10 2. The method according to claim 1, wherein the entity is an RNA:DNA hybrid-specific antibody.
3. The method according to claim 1, wherein the entity is a fragment of an RNA:DNA hybrid-specific antibody.
- 15 4. The method according to claims 2 and 3, wherein the antibody is monoclonal.
5. The method according to claims 2 and 3, wherein the antibody is polyclonal.
6. The method according to claim 1, wherein the first biological molecule is a detectably labeled double-stranded hybrid which binds the immobilized second biological molecule.
- 20 7. The method according to claim 6, wherein the immobilized second biological molecule is a protein.
8. The method according to claim 1, wherein the immobilized biological molecule comprises a plurality of biological molecules bound to a solid phase.
- 25 9. A microarray method for measuring an RNA:DNA hybrid comprising the steps of:
 - a) hybridizing a first biological molecule of the sample to a second immobilized biological molecule to form a double-stranded hybrid;
 - 30 b) measuring the amount of biological molecule of the sample; and
 - c) detecting the RNA:DNA hybrid using a detectable entity which specifically binds to a component of the RNA:DNA hybrid.
10. The method according to claim 9, wherein the entity is an RNA:DNA hybrid-specific antibody.
- 35

11. The method according to claim 9, wherein the entity is a fragment of an RNA:DNA hybrid-specific antibody.
12. The method according to claims 10 and 11, wherein the antibody is monoclonal.
- 5 13. The method according to claims 10 and 11, wherein the antibody is polyclonal.
14. The method according to claim 9, wherein the first biological molecule is a detectably labeled double-stranded hybrid which binds the immobilized second biological molecule.
- 10 15. The method according to claim 14, wherein the immobilized second biological molecule is a protein.
16. The method according to claim 9, wherein the immobilized biological molecule comprises a plurality of biological molecules bound to a solid phase.
- 15 17. A microarray method for the detection of an RNA:DNA hybrid comprising the steps of:
- a) hybridizing a first biological molecule of a sample to an immobilized second complementary biological molecule to form a double-stranded hybrid;
 - 20 b) extending the biological molecule sequence using an enzyme in conditions which promote reverse transcription; and
 - c) detecting the RNA:DNA hybrid using a detectable entity which specifically binds to a component of the RNA:DNA hybrid.
- 25 18. The method according to claim 17, wherein the entity is a detectable RNA:DNA hybrid-specific antibody.
19. The method according to claim 17, wherein the entity is a fragment of an RNA:DNA hybrid-specific antibody.
- 30 20. The method according to claims 18 and 19, wherein the antibody is monoclonal.
21. The method according to claims 18 and 19, wherein the antibody is polyclonal.
22. The method according to claim 17, wherein the immobilized biological molecule is a known expressed sequence tag (EST).
- 35

23. The method according to claim 17, wherein the immobilized biological molecule is a universal array of all possible nucleic acid sequences of at least 5- or 6- bases.
24. The method according to claim 17, wherein the enzyme is a thermostable reverse transcriptase lacking RNase H function.
25. The method according to claim 17, wherein the immobilized biological molecule comprises a plurality of biological molecules bound to a solid phase.
26. A microarray method for detecting an RNA:DNA hybrid comprising the steps of:
- a) hybridizing a first biological molecule of a sample to a complementary second biological molecule;
 - b) hybridizing the second biological molecule to an immobilized third complementary biological molecule, wherein double-stranded hybrids comprise the molecules of steps a) and b); and
 - c) detecting the RNA:DNA hybrid using a detectable entity which specifically binds to a component of the RNA:DNA hybrid.
27. The method according to claim 26, wherein the entity is a detectable RNA:DNA hybrid-specific antibody.
28. The method according to claim 26, wherein the entity is a fragment of an RNA:DNA hybrid-specific antibody.
29. The method according to claims 27 and 28, wherein the antibody is monoclonal.
30. The method according to claims 27 and 28, wherein the antibody is polyclonal.
31. The method according to claim 26, wherein steps a) and b) occur simultaneously.
32. The method according to claim 26, wherein steps a) and b) occur sequentially.
33. The method according to claim 26, wherein the immobilized biological molecule comprises a plurality of biological molecules bound to a solid phase.
34. The method according to claim 26, wherein the second biological molecule comprises:

- 0
- a) a region complementary to the first biological molecule; and
- b) a capture sequence complementary region complementary to the immobilized third biological molecule universal capture sequence.
35. The method according to claim 26, wherein the immobilized third biological molecule comprises:
- 5
- a) a common region complementary to a common region of a detectably labeled second biological molecule; and
- b) a variable region complementary to a variable region of the first biological molecule.
- 10
36. The method according to claim 26, wherein the second biological molecule is a labeled nucleic acid n-mer which binds the common region of an immobilized third complementary biological molecule.
37. The method according to claims 35 and 36, wherein the label comprises a detectable color fluorescence label and analysis utilizes a two color detection method.
- 15
38. The method according to claim 26, wherein the first biological molecule is a nucleic acid, the second biological molecule is a complementary nucleic acid to the first biological molecule, and the immobilized third biological molecule is a protein.
- 20
39. The method according to claim 26, wherein the first biological molecule is a peptide pre-bound to a nucleic acid and binds the immobilized third biological molecule of proteins, wherein the nucleic acid pre-bound to the first biological molecule is complementary to the nucleic acid of the second biological molecule.
- 25
40. The method according to claim 26, wherein the first biological molecule is a protein pre-bound to a nucleic acid and binds the immobilized third biological molecule of proteins, wherein, the nucleic acid pre-bound to the first biological molecule is complementary to the nucleic acid of the second biological molecule.
- 30
41. A method for detecting an RNA:DNA hybrid comprising the steps of:
- 35

- 56 -

- 0
- a) hybridizing a first biological molecule to a labeled second biological molecule to form a double-stranded hybrid;
 - b) binding the labeled double-stranded hybrid to an immobilized third biological molecule; and
 - 5 c) detecting the RNA:DNA hybrid using a detectable entity which specifically binds to a component of the RNA:DNA hybrid.
42. A kit for the detection of an RNA:DNA hybrid comprising all or part thereof:
- a) a sample transport medium for stabilization of the biological sample;
 - 10 b) a biological molecule which is bound to a solid phase;
 - c) a complementary biological molecule for formation of a double-stranded hybrid complex;
 - d) an enzyme for extending nucleic acid sequences;
 - 15 e) hybridization buffers which allow for hybridization;
 - f) wash buffers for eliminating any non-hybridized nucleic acid sequence; and
 - g) means for detecting the RNA:DNA hybrids using a detectable entity which specifically binds to a component of the RNA:DNA hybrid.
- 20 43. The kit according to claim 42, wherein the entity is a detectable RNA:DNA hybrid-specific antibody.
44. The kit according to claim 42, wherein the entity is a fragment of an RNA:DNA hybrid-specific antibody.
- 25 45. The kit according to claims 43 and 44, wherein the antibody is monoclonal.
46. The kit according to claims 43 and 44, wherein the antibody is polyclonal.
47. The kit according to claim 42, wherein the immobilized biological molecule comprises a plurality of biological molecules bound to a solid phase.
- 30 48. The kit according to claim 42, wherein the enzyme is a thermostable reverse transcriptase lacking RNase H function.




Figure 1A

2/23




Figure 1B

3/23




Figure 10

4/23




Figure 1D

5/23




Figure 2A

6/23




Figure 2B

7/23




Figure 20

8/23




Figure 2D

9/23





Figure 3A

10/23



Reverse Transcriptase

Figure 3B

11/23




Figure 3C

12/23




Figure 3D

13/23




Figure 3E

14/23




Figure 4A

15/23




Figure 4B

16/23





Figure 4C

17/23



Normalize Signal from RNA (green) with Signal from DNA (red)

Figure 4D

18/23





Figure 5A

19/23



Detect with anti-RNA:DNA Antibody

Figure 5B

20/23





Figure 5C

21/23



Normalize Signal from RNA (green) with Signal from DNA (red)

Figure 5D

22/23




Figure 6A

23/23




Figure 6B

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.